

RESEARCH ARTICLE

Open Access

Breastfeeding and pain relief in full-term neonates during immunization injections: a clinical randomized trial

Maryam Modarres¹, Azam Jazayeri², Parvin Rahnama^{3*} and Ali Montazeri^{4*}

Abstract

Background: The aim of this study was to examine the effect of breastfeeding on pain relief in full-term neonates during injection of hepatitis B vaccine.

Methods: This was a randomized clinical trial. A sample of full-term neonates was randomly allocated into two groups: the experimental group and the control group. Neonates in the experimental group were breastfed two minutes before, during, and after the hepatitis B immunization and the control group were held in mothers' arms but not fed. Pain was assessed using the Douleur Aiguë du Nouveau-né (DAN) scale measuring facial expressions, limb movements and vocal expressions. The assessments were carried out after hepatitis B immunization.

Results: One hundred thirty healthy full-term neonates were studied (65 in the experimental group and 65 in the control group). Gestational age, birth weight, Apgar score and gender did not differ between the two groups. The mean total pain score as measured by the DAN scale was 3.52 (SD = 1.37) for the experimental group and it was 6.78 (SD = 1.69) for the controls indicating a significant lower pain score for the experimental group ($P < 0.001$). Also, there were significant differences for the three measures of DAN scale that are facial expressions, limb movements and vocal expression, between the two study groups ($P < 0.001$).

Conclusion: The findings confirm that breastfeeding reduces pain and is effective way for pain relief during hepatitis B vaccine injection.

Trial registration: IRCT201104166206N1

Background

Healthy neonates usually experience pain during the first week of life due to several medical procedures such as intramuscular injections and heel lancing [1]. Immunization injections are the most common source of pain in childhood [2]. Untreated pain early in life may cause harmful effects on the developing central nervous system [3] and also, it might exaggerate affective and behavioral responses during subsequent painful events [4,5]. The American Academy of Pediatrics recommended that:

'Every health care facility caring for neonates should implement an effective pain-prevention program,

which includes strategies for routinely assessing pain, minimizing the number of painful procedures performed, effectively using pharmacologic and nonpharmacologic therapies for the prevention of pain associated with routine minor procedures, and eliminating pain associated with surgery and other major procedures' [6,7].

It was suggested that pain management could be based on a "3-P" approach involving pharmacologic, physical and psychological strategies [8]. Thus, studies of pain relief in neonates still are a topic of interest in pediatrics. As such breastfeeding was found to be a safe and an effective method of pain relief in newborn babies [9-13].

Several reasons were suggested to explain why breastfeeding during painful procedures might reduce pain in neonates. These could be summarized as: maternal odor

* Correspondence: p_rahnama1381@yahoo.com; montazeri@acecr.ac.ir

³Department of Midwifery, Shahed University, Tehran, Iran

⁴Mental Health Research Group, Health Metrics Research Center, Iranian Institute for Health Sciences Research, ACECR, Tehran, Iran

Full list of author information is available at the end of the article

[14], antinociceptive effect [15] skin-to-skin contact [16], the sweet tasting of milk and the act of sucking [8].

The pain reduction methods that are used in painful procedures, including the use of oral sucrose, and pacifiers have been demonstrated [17,18]. However, these methods may interfere with a correct beginning of breastfeeding [19]. Therefore it is important to assess the analgesic effect of breastfeeding as a non-pharmacological and useful approach.

In the present study, we report the results from a randomized clinical trial that assessed the efficacy of breastfeeding for pain relief during injection of hepatitis B vaccination in term neonates.

Methods

Trial design

This was a randomized, controlled trial. The study was carried out in Mirza Kochak Khan Hospital, Tehran, Iran. The painful procedure used for this study was the hepatitis B vaccination.

Participants

A sample of breastfeeding neonates was entered into the trial. Criteria for inclusion were: full-term neonate; had Apgar score of 7 and higher at five minutes after birth; delivered by spontaneous and vaginal delivery; were exclusively breastfed; postnatal age not more than 24 hours. We excluded infants with any evidence of congenital abnormalities, medical complications, or drug exposure.

Study setting

This study was conducted in Mirza Kochak Khan Hospital, Tehran, Iran.

Intervention

The study did not start until the infant was observed to be sucking at the breast. Neonates in the experimental group were breastfed during two minutes before, during, and after hepatitis B vaccination. At the end of the second minute of breastfeeding, while the infants were still sucking, an experienced nurse performed the immunization injections.

A dose of the vaccine (0.5 ml) was drawn into an Auto-disable (AD) syringe under the aseptic technique and then administered intramuscularly to the anterior thigh at a 90 angle to the skin with a 23-G, 1-inch needle. The Hepatitis B vaccine was manufactured by the Pasteur Institute of Iran and it preserved with Thimerosal. In order to minimize variability one experienced nurse, performed immunizations. For the controls the same procedure was applied while they were held in mothers' arms but not fed. None of the infants had been breastfed for at least 30 minutes before the study procedures commenced. A trained research assistant using an electronic timer was

responsible for timing of breast feeding or holding. To standardize the procedure a pilot trial with five neonates was performed.

Outcome measure

The Douleur Aiguë du Nouveau-né (DAN) scale was used to assess pain. The scale was developed by Carbajal et al. in order to measure acute pain in newborn infants [10,20]. Scores on the scale range from 0 (no pain) to 10 (maximum pain) and measures three items: facial expressions, limb movements, and vocal expression (Table 1). We asked permission from Ricardo Carbajal to use the DAN in this study. Since several studies reported different timing for the assessment of pain [9-11], we performed a pilot study and found that behavioural changes were the most important measurements for assessing pain. Also we found that the best time for assessing pain would be 45 seconds from needle injection. Scoring was performed in real time. There was an observer (outcome assessor) to complete the outcome measure.

Sample size

Based on a previous study [21], a minimum sample size of 60 infants per group (at least) was estimated. A study


Table 1 The Douleur Aiguë du Nouveau-né (DAN) scale [10,20]

Pain estimation	Score
Facial expressions	
Calm	0
Snivels and alternates gentle eye opening and closing	1
Intensity of eye squeeze, brow bulge, nasolabial furrow:	
- Mild, intermittent with return to calm*	2
- Moderate†	3
- Very pronounced, continuous‡	4
Limb movements	
Calm or gentle movements	0
Intensity of pedalling, toes spread, legs tensed and pulled up, agitation of arms, withdrawal reaction:	
- Mild, intermittent with return to calm*	1
- Moderate†	2
- Very pronounced, continuous‡	3
Vocal expression	
No complaints	0
Moans briefly (for intubated child, looks anxious or uneasy)	1
Intermittent crying (for intubated child, expression of intermittent crying)	2
Long lasting crying, continuous howl (for intubated child, expression of continuous crying)	3

* Present during <1/3 of observation periods.

† Present during 1/3 to 2/3 of observation periods.

‡ Present during >2/3 of observation periods.



with such a sample size would have a power of 90% at a 0.05 significance level. However, we recruited 65 infants for each group, giving a total sample of 130 neonates.

Randomization

Neonates were randomly assigned to the study groups. A system of sealed envelopes was used for assignment of the eligible neonates (Figure 1).

Allocation concealment

The randomization code was available only to a research fellow who was not connected to the study. The code was disclosed to the researchers when the statistical analysis was completed.

Blinding

The mothers and nurses were not blind to the group assignments. However, outcome assessor (observer) did not know the purpose and hypothesis of the study and the main investigator was blind to when the statistical analysis had been completed.

Analysis

The SPSS version 16.0 was used to analyze the data. Descriptive analyses were carried out to explore the data. The chi square test used to compare categorical variables. The t test was used to compare pain scores

between two groups. A significance level of alpha 0.05 or lower was adopted for all main analysis.

Ethics

Approval for the study was obtained from the Office for Protection of Research Subjects in Tehran University of Medical Sciences. We obtained written informed consent from mother of each neonate included in the study.

Results

Neonate characteristics

In all 130 neonates were studied (65 in the experimental group and 65 in the control group). There were no significant differences between the two groups concerning perinatal characteristics including Apgar scores ($P = 0.20$), gestational age ($P = 0.08$), neonates' weight ($P = 0.$

Table 2 Perinatal characteristics of the study samples

	Control group (n = 65)	Experimental group (n = 65)	P
	Mean (SD)	Mean (SD)	
Gestational age (weeks)	39.4 (1.2)	39.1 (1.3)	0.08
Birth weight (1000 gr.)	3.59 (0.40)	3.55 (0.38)	0.50
Apgar score	8.8 (1.2)	7.0 (1.3)	0.20
Boys/girls (numbers)	30/35	29/36	0.80

08), and gender ($P = 0.86$) (Table 2). The mean time since birth when the neonates actually received their injection was 12.0 (SD=4.03) hours.

Effect of the intervention

There was significant difference in mean of facial expressions of neonates between the control and experimental groups (2.58 (SD=0.72), 1.39 (SD=0.65) respectively). ($P < 0.001$, Table 3).

The results also showed that there was significant difference between the two groups in mean of limb movements 1.92 (SD=0.69) and experimental groups 0.83 (SD=0.51). ($P < 0.001$, Table 3).

The results of this study showed that there were significant differences in mean of Vocal expression between control 2.28 (SD=0.57) and experimental groups 1.31 (SD=0.68). ($P < 0.001$, Table 3). In addition as shown in Table 3 the difference of the DAN total score between two groups was significant.

Finally to have a better insight the DAN scores for each subscales that are facial expression, limb movements and vocal expressions are presented in Table 4.

Discussion

Fortunately, women usually do breastfed their neonates in Iran. We thought this would be a good opportunity to assess whether we could confirm previous findings on the effectiveness of breastfeeding for pain relief during a hurting medical procedure such as immunization injections. The findings from this study revealed that breastfeeding was effective for pain relief in neonates during injection of hepatitis B vaccination. This study was a step forward among similar studies for two reasons. Firstly we assured if breastfeeding was real by observing sucking movement. Secondly, breastfeeding was

Table 3 Pain evaluation with the DAN scale (0–10) according to the facial expressions, limb movements and vocal expression

	Control group (n = 65)	Experimental group (n = 65)	P*
	Mean (SD)	Mean (SD)	
Facial expressions	2.58 (0.72)	1.39 (0.65)	<0.001
Limb movements	1.92 (0.69)	0.83 (0.51)	<0.001
Vocal expression	2.28 (0.57)	1.31 (0.68)	<0.001
Total DAN Score	6.78 (1.69)	3.52 (1.37)	<0.001

* Derived from t-test.

Table 4 Pain evaluation with the DAN (0–10) in details (facial expression, limb movements and vocal expressions)

Facial expressions	Scores	Control group (n = 65)		Experimental group (n = 65)	
		Frequency	%	Frequency	%
	0	0	0	0	0
	1	3	4.6	4	6.2
	2	27	41.5	34	52.3
	3	29	44.6	25	38.5
	4	6	9.2	2	3.1
Limb movements					
	0	0	0	15	23.1
	1	18	27.7	36	70.8
	2	34	52.3	4	6.2
	3	13	20	0	0
Vocal expression					
	0	0	0	7	10.8
	1	4	6.2	32	49.2
	2	39	60	25	38.5
	3	22	33.8	1	1.5

begun two minutes before and continued during and after hepatitis B vaccination.

The finding from this study was very similar to the findings from other investigations [22]. For example, studies have shown that the breastfeeding effectively reduced response to pain during minor invasive procedure in term neonates [10] and significantly decreased crying in infants receiving immunization [12].

The role of facial expressions in measuring pain has been questioned by a recent study [23]. Therefore we reported the DAN subscales in addition to the total score to evaluate the role of each separately. However, the results showed that the three measures of DAN scale (facial expressions, limb movements, and vocal expression) were lower in the experimental group as compared to the controls.

It is argued that reactions to pain by neonates are both physiological and psychological. Consistent with our study, a study revealed that the newborns who received breastfeeding demonstrated less physiologic and behavioral responses to pain in comparison with newborns held in their mother's arms and not breastfed [1]. In addition studies have shown that breastfeeding, maternal holding, and skin to skin contact reduced crying in infants receiving an immunization injection [11].

The vaccine we used in this study was preserved with Thimerosal. In general it is argued that infant pain would be higher when using different physicochemical

properties [24,25]. However, since we used the same vaccine for all the participants, thereby the possibility of bias could be ruled out.

As indicated it is interesting to know that most women (90%) do breastfed their neonates in Iran [26]. Despite such advantage unfortunately health centers do not use breastfeeding as a non-pharmacological intervention for pain relief for immunizations or other practices [27].

Limitations

This study had some limitations. For instance selection of a single center and un-blinded nature of the study might influence the results. In addition we did not assess intra-rater reliability. Thus one should be cautious in generalizing the findings from this study. However, we were fortunate that all participants met the study criteria for the inclusion. One explanation for such compliance might be attributed to the pure chance. In addition one might question, for instance, how was that all women in this study had Apgar score of 7 or higher? The reason for this was due to the fact that all women in this study had vaginal delivery and we did not include any women who underwent caesarean section and therefore we did not consider this as failure or refusal.

Conclusion

The findings from this study suggest that breastfeeding is effective in pain reduction in immunization procedure in neonates. It may be replace practices with procedures.

Competing interests

The authors declared that they have no competing interests.

Authors' contributions

MM wrote the first draft. AJ was the main investigator, designed the study, collected the data, and performed analysis. PR helped the main investigator to analyze the data. AM contributed to the analysis, critically evaluated the paper, and provided the final draft. All authors read and approved the final version of the manuscript.

Acknowledgments

The authors gratefully acknowledge the Institutional Review Board of the Tehran University of Medical Sciences for approving and supporting this project. We are especially grateful to the chief of the pediatric department and nursing staff of the maternity ward of the Mirza Kochak Khan Hospital (women hospital). We also wish to thank the participants for their cooperation.

Author details

¹PhD student of Medical Education, Lecturer of Midwifery Education, Midwifery, Nursing & Midwifery Care Research Center, Tehran University of Medical Sciences (TUMS), Tehran, Iran. ²Faculty of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran. ³Department of Midwifery, Shahed University, Tehran, Iran. ⁴Mental Health Research Group, Health Metrics Research Center, Iranian Institute for Health Sciences Research, ACECR, Tehran, Iran.

Received: 22 July 2013 Accepted: 10 September 2013
Published: 13 September 2013

References

1. Leite AM, Linhares MB, Lander J, Castral TC, dos Santos CB, Silvan Scochi CG: **Effects of breastfeeding on pain relief in full-term newborns.** *Clin J Pain* 2009, **25**:827–832.
2. Schechter NL, Zempsky WT, Cohen LL, McGrath PJ, McMurtry CM, Bright NS: **Pain reduction during pediatric immunizations: evidence-based review and recommendations.** *Pediatrics* 2007, **119**:e1184–e1198.
3. Reis EC, Roth EK, Syphan JL, Tarbell SE, Holubkov R: **Effective pain reduction for multiple immunization injections in young infants.** *Arch Pediatr Adolesc Med* 2003, **157**:708–709.
4. Taddio A, Katz J, Ilersich AL, Koren G: **Effect of neonatal circumcision on pain response during subsequent routine vaccination.** *Lancet* 1997, **349**:599–603.
5. Taddio A, Goldbach M, Ipp M, Stevens B, Koren G: **Effect of neonatal circumcision on pain response during vaccination in boys.** *Lancet* 1995, **345**:291–292.
6. American Academy of Pediatrics: **Prevention and management of pain and stress in the neonate.** *Pediatrics* 2000, **105**:454–461.
7. American Academy of Pediatrics: **Prevention and management of pain in the neonate: an update.** *Pediatrics* 2006, **118**:2231–2241.
8. Taddio A, Appleton M, Bortolussi R, Chambers C, Dubev V, Halperin S, Hanrahan A, Ipp M, Lockett D, MacDonald N, Midmer D, Mousmanis P, Palda V, Pielak K, Ridde RP, Rieder M, Scott J, Shah V: **Reducing the pain of childhood vaccination: an evidence based clinical practice guideline (summary).** *CMAJ* 2010, **182**:1989–1995.
9. Gartner LM, Morton J, Lawrence RA, Naylor AJ, O'Hare D, Schanler RJ, Eidelman AI, American Academy of Pediatrics Section on Breastfeeding: **Breastfeeding and the use of human milk.** *Pediatrics* 2005, **115**:496–506.
10. Carbajal R, Veerapen S, Couderc S, Jugie M, Ville Y: **Analgesic effect of breast feeding in term neonates: randomized controlled trial.** *BMJ* 2003, **326**:13.
11. Efe E, Ozer ZC: **The use of breastfeeding for pain relief during neonatal immunization injections.** *Appl Nurs Res* 2007, **20**:10–16.
12. Razek AA, El-Dein NAZ: **Effect of breastfeeding on pain relief during infant immunization injections.** *Int journal of Nurs Pract* 2009, **15**:99–104.
13. Modarres M, Vasegh Rahimiparvar SF, Mehran A, Jazayeri A: **The effects of breastfeeding on pain of injection in newborns.** *Hayat* 2007, **12**:31–37 [in Persian].
14. Varendi H, Christensson K, Porter RH, Winberg J: **Soothing effect of amniotic fluid smell in newborn infants.** *Early Hum Dev* 1998, **51**:47–55.
15. Gray L, Miller LW, Philipp BL, Blass EM: **Breastfeeding is analgesic in healthy newborns.** *Pediatrics* 2002, **109**:590–593.
16. Gray L, Watt L, Blass EM: **Skin to skin contact is analgesic in healthy newborns.** *Pediatrics* 2000, **105**:e14.
17. Curtis SJ, Jou H, Ali S, Bandermeer B, Klassen T: **A randomized controlled trial of sucrose and/or pacifier as analgesia for infants receiving venipuncture in a pediatric emergency department.** *BMC Pediatr* 2007, **7**:27.
18. Blass EM, Hoffmeyer LB: **Sucrose as an analgesic for newborn infants.** *Pediatrics* 1991, **87**:215–218.
19. Howard CR, Howard FM, Lanphear B, Eberly S, deBlicek EA, Oakes D, Lawrence RA: **Randomized clinical trial of pacifier use and bottle-feeding or cup feeding and their effect on breastfeeding.** *Pediatrics* 2003, **111**:511–518.
20. Carbajal R, Paupe A, Hoenn E, Lenclen R, OlivierMartin M: **DAN: evaluation behavioral scale of acute pain in newborn infants.** *Archives Pediatre* 1977, **4**:623–628 [Article in French].
21. Blass EM, Watt LB: **Suckling- and sucrose-induced analgesia in human newborns.** *Pain* 1999, **83**:611–623.
22. Sahebshah MH, Hosseinzadeh M, Mohammadpourasl A, Kosha A: **The effect of breastfeeding, oral sucrose and combination of oral sucrose and breastfeeding in infant's pain relief during the vaccination.** *Iran J Nurs Midwifery Re* 2011, **16**:9–16.
23. Milesi C, Cambonie G, Jacquot A, Barbotte E, Mesnage R, Masson F, Pidoux O, Ferragu F, Thevenot P, Mariette JB, Picaud JC: **Validation of a neonatal pain scale adapted to the new practices in caring for preterm newborns.** *Arch Dis Child Fetal Neonatal Ed* 2010, **95**:F263–F266.
24. Tsuji Y, Ichihashi I, Morita K, Itabashi K: **Difference of the pain during the DPT (Diphtheria-Pertussis-Tetanus) vaccination.** *World J Vaccines* 2012, **2**:91–95.

25. Terada K, Inoue M, Yamaguchi T: **Difference of response in an inoculation site and pain between vaccines with Thimerosal or Phenoxyethanol.** *Infection and Immunity in Childhood* 2010, **22**:145–150.
26. Olang B, Farivar K, Heidarzadeh A, Strandvik B, Yngve A: **Breastfeeding in Iran: prevalence, duration and current recommendations.** *Int Breastfeed J* 2009, **4**:8.
27. Taavoni S, Shah-Ali SH, Neisani-Samani L, Haghani H: **Comparative study of the effect of being in mother's hug and routine clinical procedure on neonates' pain during immunization injection in health centers of west Tehran.** *Iran J Nurs* 2010, **22**:48–55.

doi:10.1186/1471-2253-13-22

Cite this article as: Modarres et al.: Breastfeeding and pain relief in full-term neonates during immunization injections: a clinical randomized trial. *BMC Anesthesiology* 2013 **13**:22.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed** Central